


"Kiney"



Open Notebook Science

Effect of Osmotic Stress and D₂O on Kinesin Activity


Andy Maloney, Brigette Black, Anthony Salvagno, Larry Herskowitz,
Brian Josey, Steve Koch, University of New Mexico



This work was supported by the DTRA CB Basic Research Program under Grant No. HDTRA1-09-1-008.
We would also like to thank Susan Atlas, PI of the DTRA project.

Motivation

Kinesin and microtubules have been proposed as components for chemical and biological sensors. In order to fully understand the dynamics of kinesin and microtubules, fundamental questions must be answered related to how we observe those interactions. One way to observe their interactions is by using a **gliding motility assay** which can easily be visualized as a molecular "crowd surfing" for microtubules. Visualizing the steps kinesin take along microtubules via fluorescence microscopy show that kinesin's motility (the ability of kinesin to walk along microtubules) is strongly affected by water's chemical potential. Changing the osmotic pressure of the aqueous environment the kinesin and microtubules are in, affects the speed at which kinesin walks along the microtubules.







Using kinesin and microtubules as chemical and biological sensors necessitates investigating how kinesin's motility is affected by osmotic stressors. Initial studies with deuterium oxide (D₂O or heavy water) have shown that kinesin's motility is affected by solvent properties. We have shown that changing the fraction of D₂O to H₂O does affect kinesin's velocity. This is a first step into understanding how we can use kinesin as a molecular reporter for chemical and biological sensors. It also introduces a new experimental "knob" if you will that will allow us to investigate the role of water on kinesin's stepping abilities.

D₂O facts



- Hygroscopic.
- 11% denser than H₂O.
- Toxic to eukaryotes. The toxic effects are similar to chemotherapeutic drugs¹.
- A mixture of 50% D₂O and 50% H₂O will dissociate to 50% HDO and 25% H₂O and 25% D₂O⁹.
- D₂O is used to stabilize viral vaccines¹⁰.
- D₂O stabilizes tubulin and microtubules².
- D₂O stimulates tubulin assembly formation².
- No previous work with kinesin has been found.

Microtubule shape differences

The differences between heavy water and light water on microtubules are stark. Some interesting shape differences include: circles, sticking, and squiggles.







All gliding motility assay images above are pseudo colored with ImageJ. Microtubules are fluorescently tagged with Rhodamine 6G.





The greener the color means more fluorescence.

Fluorescence stability

Another amazing fact about heavy water and fluorophores is that heavy water will prolong fluorescence⁴. Prolonging fluorescence actually helps the lifetime of microtubules. Photodegradation of attached fluorophores on tubulin, aid in microtubule depolymerization.



A. Polymerization in light water with 8% DMSO.
B. Polymerization in heavy water without DMSO.
C. Polymerization in heavy water with DMSO.
D. Polymerization in both heavy water and light water with 8% DMSO².




We do not polymerize our microtubules with DMSO instead, we use glycerol.


Chakrabarti et al².

Kinesin velocity


We have a joint DTRA project with a theoretical physics group lead by the PI, Susan Atlas from UNM. We are developing a new charge-transfer force field for atomistic modeling of biological systems. As a first step in our experimental tests of their new force field, we have set out to gain fundamental understanding of the effects of water activity on the enzymatic activity of kinesin. We plan to do this using osmotic stress and solvent isotope effects. Here we report our first results for the change in gliding motility assays as a function of substitution of D₂O for H₂O.



Velocity differences are not due to microtubule length. Microtubules were polymerized in light water.



Velocity dependence on D₂O concentration. Microtubules were polymerized in light water.



Microtubule length does not depend on D₂O concentration. Microtubules were polymerized in light water.

Microtubule & kinesin stability

Combined with the new and interesting features microtubules have in heavy water, they are also more stable¹. Slides last for 2 days with motility ending somewhere between day 1 and day 2. As a reference point, motility only lasts for a few hours in light water.

It has been shown that tubulin is also more stable if packed in heavy water^{2, 3}.



Microtubule polymerization

Microtubules can be polymerized in heavy water. The background fluorescence is larger in the heavy water polymerized microtubules than the light water polymerized ones. This is thought to come from ribbon structures of tubulin forming².

Future work

Continued investigation of how kinesin is affected by solvents and osmolytes will include studies that incorporate:

- **Betaines.** Betaines are zwitterionic osmolytes. They will change the osmotic pressure of the solution kinesin is in without interacting with it.
- **H₂¹⁸O.** This is a form of "heavy water" except that there is an isotope of oxygen instead of hydrogen. This may lead to interesting physics since we no longer deal with hydrogen deuterium exchange on proteins.




• **Optical tweezers.** All experiments thus far have been executed in a light microscope using gliding motility assays. The next step is to use an optical trap (designed and built by Maloney and Salvagno) to investigate kinesin attached to a dielectric bead. The optical trap will track the kinesin's motion along a microtubule thus giving us single molecule resolution of kinesin's processivity.

For more information about the experiments, please see Maloney's open notebook at: http://www.openwetware.org/wiki/User:Andy_Maloney


Microtubules

- Heterodimer of tubulin subunits α and β . One α and one β subunit together is called tubulin
- Tubulin forms polymers called protofilaments.
- Microtubules are made from 13 - 17 protofilaments.
- They are hollow and are an average of 25 nm in width.
- Calcium causes depolymerization.



Kinesin

- Dimer that consists of two heavy chains and two light chains.
- The heavy chains form the "head" group or the motor domains.
- The light chains form the "tail" group where cargo binds. The kinesin supplied to us from Dr. Liu does not have light chains. It is a truncated heavy chain Drosophila kinesin-1.
- The chains are connected by a "neck linker" and an intertwined stalk region.
- Uses ATP to generate motion.



Kinetic cycle of kinesin

Kinesin uses ATP to propel itself forward along a microtubule. The exact steps kinesin takes to complete this cycle is still debated. However, it is thought that the motor not attached to the microtubule will not rebound to it, to take a step, until an ATP binds to the motor attached to the microtubule. Binding and unbinding of kinesin motor domains to microtubules will be affected by osmotic stress and isotope water substitutions. These are experimental knobs we wish to turn in order to understand more fundamentally the kinesin catalytic cycle.

Acknowledgments

- This work was supported by the DTRA CB Basic Research Program under Grant No. HDTRA1-09-1-008.
- Haiqing Liu (CINT)
- Evan Evans
- CINT⁶ is a user facility partnered with Los Alamos and Sandia National Laboratories.
- Partial funding from the IGERT INCBN.

References

1. Panda, Dulal, Gopal Chakrabarti, Jon Hudson, Karli Pigg, Herbert P. Miller, Leslie Wilson, and Richard H. Himes. "Suppression of Microtubule Dynamic Instability and Treadmilling by Deuterium Oxide†." *Biochemistry* 39, no. 17 (May 1, 2000): 5075-5081. doi:doi: 10.1021/bi992217f.
 2. Chakrabarti, Gopal, Shane Kim, Gupta, Janice S. Barton, and Richard H. Himes. "Stabilization of Tubulin by Deuterium Oxide†." *Biochemistry* 38, no. 10 (March 1, 1999): 3067-3072. doi:doi: 10.1021/bi982461r.
 3. Das, Amlan, Sharmistha Sinha, Bipul R Acharya, Pinaki Paul, Bhabatarak Bhattacharyya, and Gopal Chakrabarti. "Deuterium oxide stabilizes conformation of tubulin:a biophysical and biochemical study." *BMB Reports* 41, no. 1 (January 31, 2008): 62-67.
 4. Sinha, S., A.K. Ray, S. Kundu, S. Sasikumar, and K. Dasgupta. "Heavy-water-based solutions of rhodamine dyes: photophysical properties and laser operation." *Applied Physics B: Lasers and Optics* 75, no. 1 (July 15, 2002): 85-90. doi:10.1007/s00340-002-0932-6.
 5. HHMI Report 2005.
 6. CINT is a user facility. More information can be found here, <http://cint.lanl.gov/>
 7. Guydosh, Nicholas R, and Steven M Block. "Direct observation of the binding state of the kinesin head to the microtubule." *Nature* 461, no. 7260 (September 3, 2009): 125-128. doi:10.1038/nature08259. Supplemental information.
 8. Klumpp, Lisa M., Andreas Hoenger, and Susan P. Gilbert. "Kinesin's second step." *Proceedings of the National Academy of Sciences of the United States of America* 101, no. 10 (March 9, 2004): 3444-3449. doi:10.1073/pnas.0307691101.
 9. http://en.wikipedia.org/wiki/Heavy_water
 10. Wu, Rong, Maria-Magdalena Georgescu, Francis Delpyroux, Sophie Guillot, Jean Balanant, Karl Simpson, and Radu Crainic. "Thermostabilization of live virus vaccines by heavy water (D2O)." *Vaccine* 13, no. 12 (1995): 1058-1063. doi:10.1016/0264-410X(95)00068-C.
- * Images false colored with ImageJ.